

INSECTICIDE

APHIDS

Registration No. L8500 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

KOMBAT APHIDS is a Suspension concentrate stomach and contact insecticide for the control of Harvest Termites, Lawn Caterpillars, Aphids, Cockroaches and various insect pests in the Home and Gardens.

Registration Holder:
KOMBAT (Pty) Ltd
Reg. No. 2009/023766/07
39 Dr Gordon Street
Greytown, 3250
Tel: +27 (0) 33 417 1906/7

Distributed by:
Starke Ayres (Pty) Ltd
Reg. No. 1984/011013/07
9th Road, Kempton Park, 1619
Tel: +27 (0) 11 748 3550
info@starkeayres.com

Active Ingredient:

Deltamethrin (pyrethroid)10,0 g/ℓ



WARNING

HAZARD STATEMENT

- May cause an allergic skin reaction.
- Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENT

- Do not breathe dust/fume/gas/mist/vapours/spray.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- Wash hands thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Avoid release to the environment.
- Collect spillage.
- Dispose of contents/container to local and national regulations.
- Call a POISON CENTER/doctor if you feel unwell.

SYMPTOMS OF POISONING

Ingesting large volumes of deltamethrin may cause nausea, vomiting, abdominal pain, and muscle twitches.

FIRST AID TREATMENT

General information: When in doubt or if symptoms are observed, get medical advice. Call a POISON CENTER/doctor if you feel unwell.

Following inhalation: No special measures are necessary. Provide fresh air.

Following skin contact: Wash with soap and water.

Following eye contact: In case of eye irritation consult an ophthalmologist.

Rinse immediately carefully and thoroughly with eye-bath or water.

Following ingestion: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Seek medical attention.

RELEVANT SUBSTANCE		CONCENTRATION (%)
Deltamethrin		
CAS N°	52918-63-5	C ≤ 1.0 %
Proxel GXL		
CAS N°	N/A	C ≤ 0.1 %

NOTE TO PHYSICIAN
Treat symptomatically.

PLEASE READ THE LABEL BEFORE USE

DIRECTIONS FOR USE

Use only as indicated.

Please note:

- Do not inhale spray mist.
- Wear gloves when preparing the spray mixture.
- Wash clothing if contaminated.
- Wash immediately after accidental skin contact with soap and cold water.
- Do not eat, drink or smoke whilst applying, mixing of before washing hands and face. Prevent contamination of fishponds, food, eating utensils and drinking water.
- Clean applicator thoroughly before re-using with other material.
- Dispose of wash water where it will not contaminate edible crops.

Rinse empty container three times with a volume of water equal to 10% of that of the container. Add the rinsing to the contents of the spray tank. Destroy empty container and do not re-use for any purpose.

1. HOMES & PUBLIC HEALTH:

ANTS: 20-80 ml per Liter water mixture applied at a rate of 50 ml/ m² as a coarse spray.

Repeat if ants re-establish themselves again.

COCKROACHES: 20 ml per Liter water mixture applied at a rate of 50 ml/ m² as a coarse spray.

2. HOME GARDENS:

NORTHERN HARVESTER TERMITE: 20 ml per 5 Liter water and apply to lawns at 5 Liter spray mixture per 100 m² to stop activity for several weeks. Repeat when activity is noticed again.

LAWN CATERPILLAR: 5 ml per 10 Liter water. Apply to lawns by means of a watering can at a rate of 10 Liter spray mixture per 20 m² when infestation is noticed.

ANTS: 5 ml per 5 Liter water mixture poured into ant nests, ensuring that the soil is well wetted. Repeat if ants re-establish themselves again.

CUTWORMS: 5 ml per 10 Liter water to treat 40 m² area. Spray as a surface spray on damp soil prior to or just after planting. A light sprinkling of water after spraying is beneficial.

CUSTOMER SERVICES: 0860 782 753 • WWW.KOMBAT.CO.ZA • MEMBER OF THE PLENNEGY GROUP

CONTROLS AND PROTECTS

APHIDS

Registration No. L8500 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

3. FLOWERS AND ORNAMENTALS:

AMERICAN BOLLWORMS: 5 mℓ / 10 Liter water. Spray at first signs of infestation. Ensure good coverage. Repeat when necessary.

APHIDS: 10 mℓ / 10 Liter water. Spray at first signs of infestation. Ensure good above and below coverage of leaves. Repeat when necessary.

AUSTRALIAN BUG: 20 mℓ / 10 Liter water. Spray when first signs of infestation are noticed. Direct spray at the insects using as high a pressure as possible to blow the egg sacs apart. Wet plant thoroughly.

CHAFER BEETLES: 5 mℓ / 10 Liter water. Spray when first leaf damage is noticed. Roses and holly-hock are good indicator plants. Beetles feed at night. Repeat when signs of feeding are seen again on new leaves.

FLOWER BEETLES (ASTYLUS BEETLES): 10 mℓ / 10 Liter water. Apply spray when beetles are first noticed, ensuring that flowers are well wetted. Roses and cannas are especially prone to attack. Repeat at 7-day intervals for as long as necessary.

SHORTHORNED GRASSHOPPERS: 15 mℓ / 10 Liter water. Spray at first sign of infestation. Repeat when necessary.

BLACK AND YELLOW BANDED LILY BORDER: 10 mℓ / 10 Liter water. Spray at first sign of infestation. Thoroughly wet the crown as larvae hide and feed there by day. Repeat spray 7 days after if reinfestation occur. Daffodils, Amaryllis and Agapanthus as well as many other bulbs are susceptible to attack.

THRIPS: 10 mℓ / 10 Liter water Gladioli leaves assume a reddish-brown speckled appearance when attack is severe. Pale marks appear on flowers which may not develop. Other plants are also prone to thrips attack, e.g. Calla, Carnation and Daffodil showing similar symptoms which should not be confused with a spider mite attack.

SHIELDBUGS: 10 mℓ / 10 Liter water. Ensure full coverage of growing tips and young growth. Repeat when necessary.

WHITE FLY: 10 mℓ / 10 Liter water. Spray at first signs of infestation. Preferably before plants become sticky with honeydew or covered in sooty mould. Repeat at weekly intervals. Ensure good coverage of both upper and lower leaf surfaces.

DISCLAIMER

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and the effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.