

INSECTICIDE

FRUITFLY

Registration No. L3343 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

KOMBAT FRUITFLY is an emulsifiable concentrate contact and stomach poison insecticide for the control of home garden pests as listed.

Active Ingredient:

Cypermethrin (pyrethroid) – 20 g/ℓ

Registration Holder:

KOMBAT (Pty) Ltd
Reg. No. 2009/023766/07
39 Dr Gordon Street
Greytown, 3250
Tel: +27 (0) 33 417 1906/7

Distributed by:

Starke Ayres (Pty) Ltd
Reg. No. 1984/011013/07
9th Road, Kempton Park, 1619
Tel: +27 (0) 11 748 3550
info@starkeayres.com

	HAZARD STATEMENT	PRECAUTIONARY STATEMENT
 WARNING	<ul style="list-style-type: none">• May cause respiratory irritation.• Very toxic to aquatic life with long lasting effects.	<ul style="list-style-type: none">• Do not breathe dust.• Wash hands thoroughly after handling.• Do not eat, drink or smoke when using this product.• Avoid release to the environment.• Wear protective gloves/ protective clothing.• Call a POISON CENTER/doctor if you feel unwell.• Collect spillage.• Dispose of contents/container to local and national regulations.

FIRST AID TREATMENT

General information:

When in doubt or if symptoms are observed, get medical advice. Call a POISON CENTER/doctor if you feel unwell.

Following inhalation: IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER/ doctor.

Following skin contact: IF ON SKIN: Wash with plenty of water. Call a POISON CENTER/doctor if you feel unwell.

Following eye contact: In case of eye irritation consult an ophthalmologist. Rinse immediately carefully and thoroughly with eye-bath or water.

Following ingestion: IF SWALLOWED: Rinse mouth.

Do NOT induce vomiting. Seek medical attention. IF SWALLOWED: Immediately call a POISON CENTER/doctor.

RELEVANT SUBSTANCE	CONCENTRATION (%)
Cypermethrin Technical (93%) (Pyrethroid)	
CAS N°	52315-07-8
	C _≤ 2.0 %

NOTE TO PHYSICIAN

Treat symptomatically.

PLEASE READ THE LABEL BEFORE USE

DIRECTIONS FOR USE

Use only as indicated.

Compatibility: Do not mix with oils, seaweed extracts and other pesticides.

Mixing and spraying procedures: To obtain better emulsion stability it is recommended to first mix the concentrate with a small quantity of water (10 ml product to 1 litre of water).

Mix thoroughly and add the rest of water until the desired volume spray is obtained.

WARNINGS

The following withholding periods must be allowed between last application and harvest:

APPLES AND PEARS	14 DAYS
BEANS AND PEA	7 DAYS
CRUCIFERAE	4 DAYS
MAIZE AND SWEETCORN	14 DAYS
PEACHES (FRESH)	14 DAYS
PEACHES (CANNING)	3 DAYS
TOMATOES	4 DAYS

DISCLAIMER

Although this remedy has been intensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or the occurrence of conditions which could not have been foreseen in the terms of the registration. Consult the supplier in the event of any uncertainty.

Please note:

- Toxic to fish and bees and moderately toxic to wildlife.
- Store in a cool place away from food and feeds.
- Keep out of reach of children and pets.
- Destroy empty container and never re-use for any other purpose.

CUSTOMER SERVICES: 0860 782 753 • WWW.KOMBAT.CO.ZA • MEMBER OF THE PLENNEGY GROUP

CONTROLS AND PROTECTS

FRUITFLY

Registration No. L3343 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

CROP	PEST	DOSAGE 10 LITRES OF WATER	REMARKS
Peaches	Fruit flies	10 mℓ	All cultivars: Apply as a full cover spray 8 weeks before picking and repeat application every 14 days. In the summer rainfall regions, the first application must be made not later than the third week of December. Peaches destined for canning can be sprayed up to 3 days before canning.
	Codling Moth	10 mℓ	Apply the first application at 75% petal drop. Repeat sprays every 14 days for high infestations. At low infestations repeat sprays every 21 days.
	False Codling Moth	10 mℓ	Early cultivars: apply as a full cover spray 6 weeks before the picking date and repeat application every 14 days.
	American Bollworm	10 mℓ	Late cultivars: apply as a full cover spray 8 weeks before the picking date and repeat every 14 days. In the summer rainfall regions the first application on must be made not later than the third week of December.
	Snoutbeetles	10 mℓ	Canning Peaches: as above but application can be made up to 3 days before picking. Apply as a full cover corrective spray when pest is noticed. Do not apply before 75% petal drop. Apply at the first signs feeding damage. Repeat application 3-4 weeks later if necessary.
All young plants	Cutworms	50 mℓ	The soil should be moist to the surface at the time of application. Apply to the soil after emergence of the plants as soon as the pest occurs. Apply at least 3ℓ of spray mixture per 100 m ² . Repeat if necessary.
Apples & Pears	Snout beetle	10 mℓ	Apply two sprays: the first at 75% petal drop and the second 4 weeks later a third spray can be applied in mid January if necessary to prevent late season damage.
	Codling Moth Leaf roller	10 mℓ	Apply the first spray at 75% petal drop, with a high leaf roller infection. Repeat every 14 days. Repeat every 21 days where a low infestation occurs.
	American Bollworm	10 mℓ	Apply as soon as larvae are noticed. Do not apply before 75% petal drop.
Beans, Peas, Maize, Sweetcorn	American Bollworm	10 mℓ	Apply as soon as larvae are noticed. Repeat every 14 days if necessary.
Cruciferae	American Bollworm Diamond Back Moth Larvae Thrips	10 mℓ	Ensure good wetting of plants. Commence spraying as soon as the first signs of the pest occur. Repeat sprays every 10-14 days.
Tomatoes	American Bollworm	10 mℓ	Apply as a full cover spray at the first signs of infestation. Repeat sprays if necessary, every 7-14 days.