

# INSECTICIDE

# KARBAPET

Registration No. L8611 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

**KOMBAT KARBAPET** is a dustable powder, with residual contact - and stomach poison, for the control of insects on various crops as listed. Also contact remedy for the control of red mites, tampans and lice on poultry, and cage birds and in poultry houses. It will also control ticks and fleas on dogs and cats.

**Active Ingredient:**


Carbaryl (carbamate) – 50 g/kg

**Registration Holder:**

KOMBAT (Pty) Ltd  
Reg. No. 2009/023766/07  
39 Dr Gordon Street  
Greytown, 3250  
Tel: +27 (0) 33 417 1906/7

**Distributed by:**

Starke Ayres (Pty) Ltd  
Reg. No. 1984/011013/07  
9th Road, Kempton Park, 1619  
Tel: +27 (0) 11 748 3550  
info@starkeayres.com

	HAZARD STATEMENT	PRECAUTIONARY STATEMENT
 <b>WARNING</b>	<ul style="list-style-type: none"> <li>Suspected of causing cancer.</li> <li>Very toxic to aquatic life.</li> </ul>	<ul style="list-style-type: none"> <li>Do not handle until all safety precautions have been read and understood.</li> <li>Avoid release into the environment.</li> <li>Wear impervious rubber gloves and boots, protective clothing and chemical safety goggles.</li> <li>If exposed or concerned: Get medical attention.</li> <li>Collect spillage.</li> <li>Store locked up.</li> <li>Dispose of contents/container to suitable landfill in accordance with local regulations.</li> </ul>

**SYMPTOMS OF POISONING**

Diarrhoea, nausea, vomiting, abdominal pains, sweating, salivation and blurred vision, tremors, muscle twitching, headache.

**FIRST AID TREATMENT**

**General information:** When in doubt or if symptoms are observed, get medical advice. Call a POISON CENTER/doctor if you feel unwell.

**Following inhalation:** IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER/ doctor.

**Following skin contact:** IF ON SKIN: Wash with plenty of water. Call a POISON CENTER/doctor if you feel unwell.

**Following eye contact:** In case of eye irritation consult an ophthalmologist. Rinse immediately carefully and thoroughly with eye-bath or water.

**Following ingestion:** IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Seek medical attention.

RELEVANT SUBSTANCE	CONCENTRATION (%)
<b>Carbaryl</b>	
CAS N°	63-25-2
	C <sub>5</sub> 5 %

**NOTE TO PHYSICIAN**

Treat symptomatically. Establish clear airway and tissue oxygenation. Administer Atropine Sulphate intravenously, or intramuscularly if iv. injection is not possible. Do not give Morphine, Aminophylline, Phenothiazines, Reserpine, Furosemide or Ethacrymic acid.

**PLEASE READ THE LABEL BEFORE USE**

**DIRECTIONS FOR USE**

Use only as indicated.

Please note:

Do not inhale dust.

Wear gloves when applying.

Wash clothing if contaminated.

Wash immediately after accidental skin contact with soap and cold water.

Do not eat, drink or smoke whilst applying, mixing of before washing hands and face.

Prevent contamination of fishponds, food, eating utensils and drinking water.

Clean applicator thoroughly before re-using with other material.

Dispose of wash water where it will not contaminate edible crops.

- Red mites and tampans in buildings: Dust all poultry houses thoroughly and pay particular attention to the floor, new litter, perches, nests, nest hay, cracks in wood and walls.
- Lice on birds: Dust under each wing and around vent.
- Ticks and fleas on dogs and cats: Sprinkle powder lightly on the hair and rub in well. In the case of cats the powder must be brushed out afterwards. Repeat weekly or as often as necessary.

**CROP PEST DOSAGE REMARKS**

**COTTON**

Red Bollworm, Spring Bollworm, Stainers, Syagrus Beetle, Jassids

and Aphids  
15 - 20 kg/ha.  
Full cover application when pest is noticed.

**LAWNS**

Caterpillar, Crickets  
20 kg/ha (20 g/10 m<sup>2</sup>)  
Apply when damage is noticed. A damp sack left on the lawn during the night will reveal infestation of lawn caterpillar when removed in the morning. Light full cover application. Water in.

**LUCERN**

Caterpillar  
15 - 20 kg/ha  
Full cover application when pest is noticed.

TOBACCO - American Bollworm, Slug  
20 kg/ha Apply when pest is noticed. Direct application to flower head in the case of American Bollworm.

**VINES**

Mealy Bug  
10 - 30 kg/ha  
Full cover application only to prevent build-up of Mealy Bug population.

Rinse empty container three times with a volume of water equal to 10% of that of the container. Add the rinsing to the contents of the spray tank. Destroy empty container and do not re-use for any purpose.

**DISCLAIMER**

Although this remedy has been intensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or the occurrence of conditions which could not have been foreseen in the terms of the registration. Consult the supplier in the event of any uncertainty.

CUSTOMER SERVICES: 0860 782 753 • WWW.KOMBAT.CO.ZA • MEMBER OF THE PLENNEGY GROUP

# CONTROLS AND PROTECTS