

INSECTICIDE

STALKBORER GRANULES

Registration No. L2491 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

KOMBAT STALKBORER GRANULES is a ready-to-use granular insecticide for the control of Stalkborer in maize.

Registration Holder:
KOMBAT (Pty) Ltd
Reg. No. 2009/023766/07
39 Dr Gordon Street
Greytown, 3250
Tel: +27 (0) 33 417 1906/7

Distributed by:
Starke Ayres (Pty) Ltd
Reg. No. 1984/011013/07
9th Road, Kempton Park, 1619
Tel: +27 (0) 11 748 3550
info@starkeayres.com

Active Ingredient:

Carbaryl (Carbamate) – 25 g/kg



WARNING

HAZARD STATEMENT	PRECAUTIONARY STATEMENT
<ul style="list-style-type: none"> Suspected of causing cancer. Very toxic to aquatic life. 	<ul style="list-style-type: none"> Do not breathe dust/fume/gas/mist/vapours/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves / protective clothing. Collect spillage. Dispose of contents/container to local and national regulations. Call a POISON CENTER/doctor if you feel unwell.

SYMPTOMS OF POISONING

The symptoms are of cholinesterase inhibition.

FIRST AID TREATMENT

General information: When in doubt or if symptoms are observed, get medical advice. Call a POISON CENTER/doctor if you feel unwell.

Following inhalation: IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER/ doctor.

Following skin contact: IF ON SKIN: Wash with plenty of water.Call a POISON CENTER/doctor if you feel unwell.

Following eye contact: In case of eye irritation consult an ophthalmologist. Rinse immediately carefully and thoroughly with eye-bath or water.

Following ingestion: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Seek medical attention.

RELEVANT SUBSTANCE	CONCENTRATION (%)
Carbaryl (Carbamate)	
CAS N°	63-25-2
	C _≤ 2.5 %

NOTE TO PHYSICIAN

Treat symptomatically. The symptoms are of cholinesterase inhibition.

Medical advice: Have the product container, label, or Material Safety Data Sheet with you when calling a poison control center or doctor or going for treatment. Consider taking venous blood for determination of blood cholinesterase activity (use heparin tube) Administer atropine sulfate, either by intramuscular or intravenously, dependent on severity of poisoning Specific antidotes are oximes (e.g., Pralidoxime) or Toxogonin.

PLEASE READ THE LABEL BEFORE USE

DIRECTIONS FOR USE

Use only as indicated.

ALLOW 14 DAYS BETWEEN APPLICATION AND GRAZING OR HARVEST.

Please note:

- May be harmful if swallowed.
- Toxic to fish and wildlife.
- Store away from food and feed.
- Keep out of reach of children, uninformed persons and animals.

Destroy empty container and never re-use for any other purpose.

KOMBAT STALKBORER GRANULES ARE READY TO USE.

CROP/PEST	DOSAGE	REMARKS
MAIZE: First Generation Stalkborer / mealie grub (Busseola fusca)	40 g / 100 metre plant row (approx. 4 kg/ha)	Apply as soon as the characteristic shotholes appear on the young leaves of the maize plants. Apply the granules to the funnels of the plants using a container with a perforated lid.

DISCLAIMER

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and the effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

CUSTOMER SERVICES: 0860 782 753 • WWW.KOMBAT.CO.ZA • MEMBER OF THE PLENNEGY GROUP

CONTROLS AND PROTECTS