

INSECTICIDE

TERMITES

Registration No. L4981 – ACT 36 OF 1947



Member of the PLENNEGY GROUP

KOMBAT TERMITES is a bait, in granular form, for the control of Northern harvester termites on lawns, lucern fields, wheat fields, around lucerne stores, haystacks and pastures.

Registration Holder:
KOMBAT (Pty) Ltd
Reg. No. 2009/023766/07
39 Dr Gordon Street
Greytown, 3250
Tel: +27 (0) 33 417 1906/7

Distributed by:
Starke Ayres (Pty) Ltd
Reg. No. 1984/011013/07
9th Road, Kempton Park, 1619
Tel: +27 (0) 11 748 3550
info@starkeayres.com

Active Ingredient:

Carbaryl (Carbamate) – 16 g/kg



WARNING

HAZARD STATEMENT

- Suspected of causing cancer.
- Very toxic to aquatic life.

PRECAUTIONARY STATEMENT

- Do not handle until all safety precautions have been read and understood.
- Avoid release into the environment.
- Wear impervious rubber gloves and boots, protective clothing and chemical safety goggles.
- Collect spillage.
- Store locked up.
- Dispose of contents/container to suitable landfill in accordance with local regulations.

SYMPTOMS OF POISONING

Symptoms may include nausea, vomiting, diarrhoea, abdominal cramps, headache, vertigo, ocular pain, ciliary muscle spasm, blurring or dimness of vision, miosis, or in some cases mydriasis, lacrimation, salivation, sweating, and confusion. In severe cases, there may also be involuntary defecation and urination, bradycardia, hypotension, pulmonary oedema, convulsions, coma, and death from respiratory failure or cardiac arrest. Carbaryl does not accumulate in mammalian tissue and the cholinesterase inhibition reverses rather rapidly. In non-fatal cases, the illness generally lasts less than 24 hours.

Chronic exposure: Prolonged or repeated exposure may cause effects as described in acute exposure.

FIRST AID TREATMENT

General information: When in doubt or if symptoms are observed, get medical advice. Call a POISON CENTER/doctor if you feel unwell.

Following inhalation: IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER/ doctor.

Following skin contact: IF ON SKIN: Wash with plenty of water. Call a POISON CENTER/doctor if you feel unwell.

Following eye contact: In case of eye irritation consult an ophthalmologist.

Rinse immediately carefully and thoroughly with eye-bath or water.

Following ingestion: IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Seek medical attention. IF SWALLOWED: Immediately call a POISON CENTER/doctor.

RELEVANT SUBSTANCE		CONCENTRATION (%)
Carbaryl (Carbamate)		
CAS N°	63-25-2	C _≤ 1.6 %

NOTE TO PHYSICIAN

Treat symptomatically. Treat respiratory difficulty with artificial respiration and oxygen. Do not give morphine, aminophylline, phenothiazines, reserpine, furosemide, or ethacrynic acid. Drugs like 2 PAM are not effective in poisoning with Carbaryl. THEY SHOULD NOT BE USED. Treat symptomatically and supportively. Get medical attention immediately.

PLEASE READ THE LABEL BEFORE USE

DIRECTIONS FOR USE

Use only as indicated.

Please note:

- Withdraw livestock from treated areas during and for a period of 3 days after treatment.
- Toxic to fish.
- Store in a cool place away from food and feedstuffs. Keep out of reach of children, poultry, livestock and domestic animals.
- Destroy empty container and never use for any other purpose.
- Do not eat, drink or smoke whilst applying or before having washed hands.

Ensure that the labourers or application machinery do not smell of bone meal, fish meal, tobacco, fats, oils, petrol or other strong-smelling substances. Bait will repel termites if contaminated with minute quantities of sharp smelling substances. Avoid wet conditions because rain destroys the bait.

TARGET	DOSAGE	REMARKS
Pastures, against the harvester termite <i>Hodotermes mossambicus</i> which does not build mounds.	10 kg/ha per application 1 g/m ²	Scatter around the foraging holes while the termites are intensely foraging. Do not apply more than they will carry in that day. They are most active from 11:00 to sundown, especially from May to August. Repeat 2-4 days later and again 10 days after that in case the harvesting has not stopped. As the brown adults die off, only white nymphs will come up to forage. 2-4 weeks of well-timed applications will probably destroy the Colonies, altogether. However, spot treatment may have to continue where on large areas the one or another nest has been overlooked. Avoid application during wet weather conditions.

DISCLAIMER
Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and the effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

CUSTOMER SERVICES: 0860 782 753 • WWW.KOMBAT.CO.ZA • MEMBER OF THE PLENNEGY GROUP

CONTROLS AND PROTECTS